

**CALORIMETRIC STUDY OF THE OXOTRANSFER PROCESS INVOLVING  
TRIPHENYLPHOSPHINE AND A MOLYBDENUM-CONTAINING  
ANALOGUE OF THE ENZYME ACTIVE SITE**

*Simonova V.M.*<sup>(1)</sup>, *Podturkina S.A.*<sup>(1)</sup>, *Khripun V.D.*<sup>(2)</sup>, *Pestova O.N.*<sup>(1)</sup>

<sup>(1)</sup> St. Petersburg State University

199034, Saint Petersburg, Universitetskaya emb., 7-9

<sup>(2)</sup> Federal University of Mato Grosso do Sul

79070-900, Campo Grande, Brazil, Av. Costa e Silva

The key process that forms the basis of molybdenum enzymes' operation is oxotransfer, i.e. the transfer of an oxygen atom between the enzyme's active site and the substrate. Molybdenum complexes with dithiolene ligands are commonly used as convenient models of the active site in molybdenum-containing enzymes. Research on these model active sites typically aims to obtain kinetic and electrochemical characteristics of the system. Nevertheless, thermodynamic data are essential for identifying the driving force behind the oxotransfer process [1].

In this study, the molybdenum complex with maleonitriledithiolate ligand ( $\text{mnt}^{2-}$ ) of the composition  $(\text{Bu}_4\text{N})_2[\text{MoO}_2(\text{mnt})_2]$  was used as a model of the enzyme active center, and triphenylphosphine  $\text{PPh}_3$  was used as the substrate. The calorimetric study was performed using Calvet-type microcalorimeter (KYV) [2] to obtain data on the change in enthalpy during the dissolution processes of the solid sample  $(\text{Bu}_4\text{N})_2[\text{MoO}_2(\text{mnt})_2]$  (a) in pure acetonitrile and (b) in acetonitrile containing a portion of  $\text{PPh}_3$ . Subsequently, the total changes in enthalpy upon interaction of the components in the liquid phase were calculated.

By varying the ratio of the complex to the substrate, the values of the total thermal effects during the oxotransfer process in this system were obtained. The increase in the phosphine quantity in the system leads to a significant increase in total the thermal effect. These data indicate a catalytic character of the process, which correlates with the nature of the synthetic model.

1. Simonova, V. M., Pestova, O. N., Anufrikov, Y. A., Espósito, B. P., Khripun, V. D. Oxidation of  $\text{PPh}_3$  by Oxygen Catalyzed by Biomimetic Tungsten Complex: Oxo-Transfer Process Thermodynamics. // J Therm Anal Calorim. 2024. Vol. 149, P. 4771-4781. <https://doi.org/10.1007/s10973-024-13065-z>

2. Golubev B.A., Kondratiev Yu.V., and Sheitelman B.I. (1985) USSR Inventor's Certificate no. 1283553.

*The work was performed using the equipment of the St. Petersburg State University Science Park (CMR, CAMR and XRD Centre) and was carried out as part of an initiative project supported by St. Petersburg State University, grant 149039462. VDK acknowledges funding from FUNDECT/CNPq (grant 83/053.361/2024) The authors express special gratitude to Dr. Kondratiev Yu.V., who designed the KYV calorimeter.*